

**Problem 1: Pauli matrices and Bloch sphere.**

1. Check the relation  $\sigma_\alpha \sigma_\beta = i\varepsilon_{\alpha\beta\gamma} \sigma_\gamma + \delta_{\alpha\beta} I$  for the Pauli matrices  $\sigma_\alpha$ ,  $\alpha, \beta, \gamma = 1, 2, 3$ .
2. The *trace*  $\text{tr}[X]$  is defined as the sum of the diagonal elements of  $X$ , i.e.,  $\text{tr}[X] := \sum_i X_{ii}$ . Determine  $\text{tr}[I]$ ,  $\text{tr}[\sigma_\alpha]$ , and  $\text{tr}[\sigma_\alpha \sigma_\beta]$ .
3. Determine the eigenstates (=eigenvectors) and eigenvalues of the Pauli matrices.
4. Determine the angles  $\theta$  and  $\phi$  of those eigenstates on the Bloch sphere, and depict their position on the Bloch sphere.
5. Given a state

$$|\psi\rangle = e^{i\chi} [\cos(\theta/2)|0\rangle + e^{i\phi} \sin(\theta/2)|1\rangle] \quad (1)$$

show that

$$|\psi\rangle\langle\psi| = \frac{1}{2}(I + \vec{v} \cdot \vec{\sigma}) \quad \text{with } \vec{v} \in \mathbb{R}^3 \text{ and } |\vec{v}| = 1, \quad (2)$$

(i.e.,  $\vec{v}$  is a vector on the unit sphere in  $\mathbb{R}^3$ ), where  $\vec{v} \cdot \vec{\sigma} = \sum_{i=1}^3 v_i \sigma_i$ . (You should find that  $\vec{v}$  is exactly the point on the Bloch sphere with spherical coordinates in  $\theta$  and  $\phi$ , just as introduced in the lecture.)

6. Show that the expectation value of the Pauli operators is  $\langle\psi|\sigma_i|\psi\rangle = v_i$ ; i.e.,  $|\psi\rangle$  describes a spin which is polarized along the direction  $\vec{v}$ .
7. Show that for any state  $|\psi\rangle$  with corresponding Bloch vector  $\vec{v}$ , the state  $|\phi\rangle$  orthogonal to it, i.e. with  $\langle\psi|\phi\rangle = 0$  (for qubits, i.e., in  $\mathbb{C}^2$ , this state is uniquely determined up to a phase!), is described by the Bloch vector  $-\vec{v}$ , i.e., it is located at the opposite point of the Bloch sphere.  
(*Bonus question:* Derive a general expression for the overlap  $|\langle\phi|\psi\rangle|^2$  of two arbitrary states in terms of the corresponding Bloch vectors.)

(*Note:* A particularly elegant way to check 6. and 7. is to use that  $\langle\psi|O|\psi\rangle = \text{tr}[|\psi\rangle\langle\psi|O]$  – this is easily shown by writing this explicitly as a sum over components, but you can just this formula as it is if you want, as it will be proven in the lecture on Monday, 19.10. – together with Eq. (2) and  $\text{tr}[\sigma_i \sigma_j] = 2\delta_{ij}$ , but the results can of course also be derived directly from Eq. (1) with a bit more brute force.)

**Problem 2: Matrix spaces as Hilbert spaces.**

Let  $\mathcal{V}_d$  be the space of all complex  $d \times d$  matrices, and  $\mathcal{W}_d \subset \mathcal{V}_d$  the space of all hermitian complex  $d \times d$  matrices (i.e. for  $M \in \mathcal{W}_d$ ,  $M = M^\dagger$ ).

1. Show that  $\mathcal{V}_d$  forms a vector space over  $\mathbb{C}$ , and  $\mathcal{W}_d$  forms a vector space over  $\mathbb{R}$ , but not over  $\mathbb{C}$ . We will in the following always consider  $\mathcal{V}_d$  as a complex and  $\mathcal{W}_d$  as a real vector space.
2. Show that the Pauli matrices together with the identity,  $\Sigma := \{\sigma_i\}_{i=0}^3$ , form a basis for both  $\mathcal{V}_2$  (over  $\mathbb{C}$ ) and  $\mathcal{W}_2$  (over  $\mathbb{R}$ ).
3. Show that

$$(A, B) = \text{tr}[A^\dagger B]$$

defines a scalar product (the “Hilbert-Schmidt scalar product”) both for  $\mathcal{V}_d$  and for  $\mathcal{W}_d$ . Here,  $\text{tr}[X]$  is the trace, i.e., the sum of the diagonal elements.

4. Show that the Pauli matrices  $\Sigma$  form an orthonormal basis (ONB) with respect to the Hilbert-Schmidt scalar product.

5. Use the fact that for any scalar product  $(\vec{v}, \vec{w})$  and a corresponding ONB  $\vec{w}_i$ , we can write

$$\vec{v} = \sum_i \vec{w}_i (\vec{w}_i, \vec{v}) ,$$

to express a general matrix in  $M \in \mathcal{V}_2$  as

$$M = \sum m_i \sigma_i .$$

What is the form of the  $m_i$ ? What special property do the  $m_i$  satisfy for  $M \in \mathcal{W}_2$ ?

6. Show that a hermitian orthonormal basis also exists for  $\mathcal{V}_d$  and  $\mathcal{W}_d$ . (Ideally, explicitly construct such a basis.)

### Problem 3: Unitary invariance and Bell states.

1. Show that the singlet state

$$|\Psi^-\rangle = \frac{1}{\sqrt{2}} (|01\rangle_{AB} - |10\rangle_{AB})$$

is invariant under joint rotations by the same  $2 \times 2$  unitary  $U$ , i.e.,

$$|\Psi^-\rangle = (U \otimes U) |\Psi^-\rangle$$

for any unitary matrix  $U$ ,  $U^\dagger U = I$ .

2. Show that this implies that if we measure the spin in any direction  $\vec{v}$ ,  $|\vec{v}| = 1$  – this measurement is described by the measurement operator  $S_{\vec{v}} = \sum_{i=1}^3 v_i \sigma_i$  – we obtain perfectly random and opposite outcomes.

(*Hint:* An elegant way of doing so is to first show that any  $S_{\vec{v}}$  has the same eigenvalues as the  $Z$  matrix and therefore can be rotated to it, i.e., there exists a  $U_{\vec{v}}$  s.th.  $U_{\vec{v}} S_{\vec{v}} U_{\vec{v}}^\dagger = Z$ . Note that there are very elegant ways to show that the eigenvalues are  $\pm 1$  as well!)

3. Determine the states

$$\begin{aligned} (X \otimes I) |\Psi^-\rangle , & \quad (I \otimes X) |\Psi^-\rangle , \\ (Y \otimes I) |\Psi^-\rangle , & \quad (I \otimes Y) |\Psi^-\rangle , \\ (Z \otimes I) |\Psi^-\rangle , & \quad (I \otimes Z) |\Psi^-\rangle . \end{aligned}$$

Why are they pairwise equal?

*Note:* Together with  $|\Psi^-\rangle$ , these are known as the four *Bell states*.

4. Show that the maximally entangled state

$$|\Omega\rangle = \sum_{i=1}^d |i, i\rangle$$

of two qu- $d$ -its (i.e., systems with a Hilbert space  $\mathbb{C}^d$ ) is invariant under  $U \otimes \bar{U}$ , where  $U$  is any  $d \times d$  unitary, that is,

$$|\Omega\rangle = (U \otimes \bar{U}) |\Omega\rangle .$$