

2. Mixed States

a) The density operator

Considers a bipartite state $|\psi\rangle_{AB} = \sum c_{ij} |i\rangle |j\rangle$.

We have access to A only.

Can we characterize the measurement outcomes for meas. on A in a simple way?

(i.e. without having to consider B, which we anyway cannot access!)

Considers measurement operator Π .

(e.g. $\Pi = E_i$ projector, or exp. value, ...)

Measurement of $\Pi \equiv \Pi_A$ on A

\iff measurement of $\Pi_A \otimes I_B$ on $A \otimes B$.

$$\begin{aligned} \langle \psi | \Pi_A \otimes I_B | \psi \rangle &= \sum_{\substack{i'j' \\ ij'}} \overline{c_{i'j'}} \langle i' | \langle j' | (\Pi_A \otimes I_B) | i \rangle | j \rangle c_{ij} \\ &= \sum \overline{c_{i'j'}} c_{ij} \langle i' | \Pi_A | i \rangle \underbrace{\langle j' | j \rangle}_{= \delta_{j'j}} \end{aligned}$$

$$= \sum_{ii'} \left(\sum_j \bar{c}_{ij} c_{ij} \right) \langle i' | \rho_A | i \rangle = (*)$$

Now: Define ρ_A - a $d_A \times d_A$ matrix - via

$$(\rho_A)_{ii'} = \sum_j c_{ij} \bar{c}_{ij} = (C \cdot C^\dagger)_{ii'}$$

with the matrix $C = (c_{ij})_{ij}$,

$$\text{or equivalently } \rho_A = \sum_{i, i', j} c_{ij} \bar{c}_{ij} |i\rangle\langle i'|$$

and introduce the trace

$$\text{tr}(X) = \sum_k \langle k | X | k \rangle$$

can be any ONB,
not necessarily comp.
basis.

Note: The trace is

• cyclic: $\text{tr}(AB) = \sum_k \langle k | AB | k \rangle$

$$= \sum_k \langle k | A \left(\sum_e |e\rangle\langle e| \right) B | k \rangle$$

$$= \sum_{ke} \langle k | A | e \rangle \langle e | B | k \rangle$$

$$= \sum_{ke} \langle e | B | k \rangle \langle k | A | e \rangle = \text{tr}(BA)$$

Note: A, B

need not

be square!

- and thus basis-independent:

$$\text{tr}(u^\dagger X u) = \text{tr}(X u u^\dagger) = \text{tr}(X),$$

and thus $\text{tr}(X) = \sum \langle k | X | k \rangle$

$$= \sum \underbrace{\langle k | u^\dagger}_{\langle v_k |} X \underbrace{(u | k \rangle)}_{|v_k \rangle}$$

$$= \sum \langle v_k | X | v_k \rangle \text{ for any ONB,}$$

- the sum of the eigenvalues:

$$\text{tr}(X) = \text{tr}(X A A^\dagger) = \text{tr}(A^\dagger X A),$$

with $A^\dagger X A$ the eigenvalue decomposition.

- and of course linear:

$$\text{tr}(A) + \lambda \text{tr}(B) = \text{tr}(A + \lambda B).$$

Then,

$$\begin{aligned}
 (*) &= \sum_{ii'} \left(\sum_j \overline{c_{ij}} c_{ij} \right) \langle i' | \Pi_A | i \rangle \\
 &= \sum_{ii'} \left(\sum_j \overline{c_{ij}} c_{ij} \right) \text{tr} \left[\langle i' | \Pi_A | i \rangle \right] \\
 &= \sum_{ii'} \left(\sum_j c_{ij} \overline{c_{ij}} \right) \text{tr} \left[|i\rangle \langle i'| \Pi_A \right]
 \end{aligned}$$

trace of a number is itself!

} cyclicity of trace!

linearity of trace!

$$\downarrow = \text{tr} \left[\left(\sum_{ii'} c_{ij} \overline{c_{ij}} |i\rangle \langle i'| \right) \Pi_A \right]$$

$$= \rho_A$$

$$= \text{tr} [\rho_A \Pi_A].$$

i.e.: $\langle \psi | \Pi_A \otimes \mathbb{I}_B | \psi \rangle = \text{tr} [\rho_A \Pi_A],$

where $\rho_A = \sum_{ii'} c_{ij} \overline{c_{ij}} |i\rangle \langle i'|,$

or $\rho_A = C C^T,$ with $C = (c_{ij})_{ij}.$

ρ_A is called the density operator, density matrix,
or mixed state. It characterizes systems where
we only have partial knowledge, such as
access to only part of the system.

In contrast, a state $|\psi\rangle \in \mathcal{H}$ is called a pure state.

If we want to highlight that ρ_A comes from a larger system,
we can also refer to it as the reduced density matrix of
system A.

Properties of ρ_A :

- $\rho_A = CC^\dagger \Rightarrow \rho_A^\dagger = (CC^\dagger)^\dagger = CC^\dagger = \rho_A$

- ρ_A is positive semidefinite:

$$\begin{aligned}\langle \phi | \rho_A | \phi \rangle &= \langle \phi | CC^\dagger | \phi \rangle = (C^\dagger | \phi \rangle)^\dagger \underbrace{(C^\dagger | \phi \rangle)}_{=: |\phi'\rangle} \\ &= \langle \phi' | \phi' \rangle \geq 0 \quad \forall \phi.\end{aligned}$$

We write $\rho_A \geq 0$.

Note: $X \geq 0$, i.e. $\langle \phi | X | \phi \rangle \geq 0 \quad \forall |\phi\rangle$

$\iff X = X^\dagger$ & all eigenvalues of X are ≥ 0 .

(In part., $X \geq 0 \Rightarrow X = X^\dagger$)

Chapter II, pg 46

- $\text{tr}(\rho_A) = \sum_i (C C^\dagger)_{ii} = \sum_{ij} c_{ij} \bar{c}_{ij} = \sum |c_{ij}|^2 = 1.$

Properties of density operators:

- $\rho_A \geq 0$ (implies $\rho_A = \rho_A^\dagger$)
- $\text{tr}(\rho_A) = 1.$

Will see soon: This provides an alternative fundamental definition of a state — i.e., any ρ_A with the properties above can arise if we only have access to part of the system.

Note: All ρ_A with the above property form a convex set S , i.e.:

$$\rho, \sigma \in S \Rightarrow p\rho + (1-p)\sigma \in S, \quad 0 \leq p \leq 1.$$

Is there an ambiguity in ρ_A , just as the phase ambiguity for pure states?

Theorem: ρ_A is uniquely determined by all measurement outcomes $\text{tr}[\rho_A \Pi]$ for $\Pi = \Pi^\dagger$.

(i.e., by all overlaps, through probabilities, i.e. Π orth. proj., also suffices.)

Proof: Let $V = \{ \Pi \mid \Pi = \Pi^\dagger \}$. V is a vector space over \mathbb{R} .

$(\Pi, N) := \text{tr}[\Pi^\dagger N]$ defines a scalar product on V (the "Hilbert-Schmidt scalar product").

Pick an ONB $\{ \pi_i \}$ of V , $\text{tr}[\pi_i^\dagger \pi_j] = \delta_{ij}$.

Then, the map $X \mapsto \sum \pi_i \text{tr}[\pi_i^\dagger X]$
 $= \sum \pi_i (\pi_i, X)$

acts as the identity on V . Thus,

$$\rho_A = \sum \pi_i \text{tr}[\pi_i \rho_A],$$

i.e., ρ_A is fully specified by all meas. outcomes

(and thus, there must be a unique ρ_A for any given physical state. \square)

(Note: We didn't really use that we have hermitian matrices - the same ideas work for $V_{\mathbb{C}} = \{\pi\}$ over \mathbb{C} . Then the \cdot^+ are important - and we must show that $V_{\mathbb{C}}$ has a hermitian basis over \mathbb{C} - which it does.)

In particular: No ambiguity in P_A

\Rightarrow all numbers meaningful!

Where did the phase $|\psi_A\rangle \sim e^{i\phi} |\psi_A\rangle$ go?

Density matrix for a pure state $|\psi_A\rangle$:

$$\langle \psi_A | \pi | \psi_A \rangle = \text{tr}[\langle \psi_A | \pi | \psi_A \rangle]$$

number
↓

$$= \text{tr}[\pi \underbrace{|\psi_A\rangle\langle\psi_A|}_{= P_A}]$$

↑
cycl.

$\Rightarrow P_A = |\psi_A\rangle\langle\psi_A|$: projector onto $|\psi_A\rangle$.

(Phase naturally drops out!)


b) The partial trace

Just seen: Pure state on $AB \rightarrow$ Mixed state on A .

What if AB itself is already mixed
(e.g. from a pure ABC ?)

Same approach: How to describe most general measurement on A , given a state ρ_{AB} ?

$$\text{tr}[(\Pi_A \otimes I_B) \rho_{AB}] = \sum \langle ij | \Pi \otimes I | i'j' \rangle \langle i'j' | \rho_{AB} | ij \rangle$$



 $= \delta_{ij'}$

$$= \sum_{ii'} \langle i | \Pi | i' \rangle \langle i'j' | \rho_{AB} | ij \rangle$$

$$= \text{tr} \left[\Pi \cdot \left(\sum_{ii'} |i'\rangle_A \langle i'j' | \rho_{AB} | ij \rangle \langle i | \right) \right]$$

$$= \text{tr}[\Pi \cdot \rho_A]$$

where we define

$$\rho_A = \sum |i'\rangle_A \langle i'j' | \rho_{AB} | ij \rangle \langle i | \rho_A$$

$$\begin{aligned}
&= \sum_j (I_A \otimes \langle j|_B) P_{AB} (I_A \otimes |j\rangle_B) \\
&= \sum_j \langle j|_B P_{AB} |j\rangle_B \\
&=: \text{tr}_B(P_{AB}) : \text{the } \underline{\text{"partial trace"}}
\end{aligned}$$

In components:

$$\begin{aligned}
(\text{tr}_B(P_{AB}))_{ii'} &= \langle i|_A \left(\sum_j \langle j|_B P_{AB} |j\rangle_B \right) |i'\rangle \\
&= \sum_j (P_{AB})_{(ij), (i'j)}
\end{aligned}$$

(Note: The partial trace can also be seen as the canonical embedding of

$$\text{tr}: B(\mathcal{H}_A) \rightarrow \mathbb{C}$$

into $B(\mathcal{H}_A) \otimes B(\mathcal{H}_B)$

linear ("bounded") operators on \mathcal{H}_A .

Note: P_A is also called reduced density matrix (or operator) of P_{AB} (or ρ_{AB}).

c) Purification

Is any density matrix ρ ($\rho \geq 0, \text{tr} \rho = 1$) physical (i.e., coming from a pure state, as by our axioms)?

Purification of mixed state ρ :

Consider any decomposition $\rho = \sum \lambda_i |\phi_i\rangle\langle\phi_i|$, $\lambda_i > 0$, e.g. the eigenvalue decomposition, and define

$$|\psi\rangle_{AB} := \sum \sqrt{\lambda_i} |\phi_i\rangle_A |i\rangle_B$$

any ONB

$$\text{Then } \text{tr}_B [|\psi\rangle\langle\psi|] = \text{tr}_B \left[\sum_{ij} \sqrt{\lambda_i \lambda_j} |\phi_i\rangle\langle\phi_j| \otimes |i\rangle\langle j| \right]$$

$$= \sum_{ij} \sqrt{\lambda_i \lambda_j} |\phi_i\rangle\langle\phi_j| \otimes \underbrace{\text{tr}_B [|i\rangle\langle j|]}_{= \delta_{ij}}$$

$$= \sum \lambda_i |\phi_i\rangle\langle\phi_i| = \rho$$

Yes, every ρ is physical (in the sense above).

\Rightarrow Density operator ρ can serve as an alternative fundamental definition of a state in quantum theory.

Definition: A $|\psi\rangle_{AB}$ s.t. $\text{tr}_B(|\psi\rangle\langle\psi|) = \rho$ is called a purification of ρ .

Note: The ambiguity of purifications - i.e., how are two purifications $|\psi\rangle, |\phi\rangle$ of ρ , $\text{tr}_B(|\psi\rangle\langle\psi|) = \text{tr}_B(|\phi\rangle\langle\phi|) = \rho$, related - will be addressed later.

d) Ensemble interpretation of the density matrix

Consider $|\psi\rangle = \alpha|00\rangle + \beta|11\rangle$:

$$\Rightarrow \rho_A = \begin{pmatrix} |\alpha|^2 & 0 \\ 0 & |\beta|^2 \end{pmatrix} = |\alpha|^2 |0\rangle\langle 0| + |\beta|^2 |1\rangle\langle 1|$$

$$\Rightarrow \text{tr}[\rho_A] = |\alpha|^2 \langle 0|\rho|0\rangle + |\beta|^2 \langle 1|\rho|1\rangle.$$

\Rightarrow Can be interpreted as having the pure state $|0\rangle$ with probability $p_0 = |\alpha|^2$, and $|1\rangle$ w/ $p_1 = |\beta|^2$.

"ensemble interpretation" of density matrix

However: We have derived ρ_A from a pure state

$|\psi\rangle_{AB}$ — are these two perspectives consistent?

Imagine B does a measurement in the Z basis:

$$|\psi\rangle = \alpha|00\rangle + \beta|11\rangle$$

$p_0 = |\alpha|^2 \rightarrow |\psi_0\rangle_A = |0\rangle_A$
 $p_1 = |\beta|^2 \rightarrow |\psi_1\rangle_A = |1\rangle_A$

The post-measurement state of Alice is $|\psi_0\rangle = |0\rangle$
 with $p_0 = |\alpha|^2$, and $|\psi_1\rangle = |1\rangle$ with $p_1 = |\beta|^2$.

But: Alice does not know outcome of Bob

\Rightarrow meas. of B produces an ensemble

$$\{ (p_0, |0\rangle), (p_1, |1\rangle) \} =$$

$$= p_0 |0\rangle\langle 0| + p_1 |1\rangle\langle 1| = \begin{pmatrix} p_0 & \\ & p_1 \end{pmatrix} = \begin{pmatrix} |\alpha|^2 & \\ & |\beta|^2 \end{pmatrix}.$$

(But what: Bob knows outcome \Rightarrow his description

is different: he would describe Alice's state

either as $|0\rangle\langle 0|$ or as $|1\rangle\langle 1|$!

i.e.: state assigned dep. on knowledge!

But: Bob could also measure in different bases,

$$\text{e.g. } |\pm\rangle = \frac{1}{\sqrt{2}} (|0\rangle \pm |1\rangle)!$$

$$P_+ = \frac{|\alpha|^2 + |\beta|^2}{2} = \frac{1}{2} \rightarrow |Y_+\rangle_A = \frac{\alpha|0\rangle + \beta|1\rangle}{|\alpha|^2 + |\beta|^2}$$

$$|\psi\rangle = \alpha|00\rangle + \beta|11\rangle$$

X-meas.
on B

$$P_- = \frac{|\alpha|^2 + |\beta|^2}{2} = \frac{1}{2} \rightarrow |Y_-\rangle_A = \frac{\alpha|0\rangle - \beta|1\rangle}{|\alpha|^2 + |\beta|^2}$$

non-orthogonal!

Ensemble $\{(P_+, |Y_+\rangle), (P_-, |Y_-\rangle)\}$

Indeed, $P_+ |Y_+\rangle \langle Y_+| + P_- |Y_-\rangle \langle Y_-| = \rho_A!$

Different ensemble for same state

\Rightarrow ensemble interpretation is ambiguous!

(Even # of terms can vary, etc. \rightarrow HWS)

Definition: We call a system (or a collection of systems) which is in state $|\psi_i\rangle$ (or ρ_i) with prob. P_i an ensemble. (We write $\{(P_i, |\psi_i\rangle)\}$, or $\{(P_i, \rho_i)\}$.)

Observation: Measurement outcomes for an ensemble

$\{(\rho_i, p_i)\}$ are described by


$$\langle \Pi \rangle := \sum p_i \operatorname{tr}[\Pi \rho_i] = \operatorname{tr}[\Pi (\underbrace{\sum p_i \rho_i}_{=: \rho})]$$

↑
any.

$$= \operatorname{tr}[\Pi \rho]$$

\Rightarrow Different ensembles $\sum p_i \rho_i = \sum p'_i \rho'_i$ are indistinguishable.

How are two different ensemble decompositions related?



Theorem: $\sum_i p_i |\psi_i\rangle\langle\psi_i| = \sum_j q_j |\phi_j\rangle\langle\phi_j|$

no need for orthonormality!!

if and only if there exists $U = (u_{ij})$ s.t.

$$\sqrt{p_i} |\psi_i\rangle = \sum_j u_{ij} \sqrt{q_j} |\phi_j\rangle,$$

where $U = (u_{ij})$ satisfies

(i) $\sum_j q_j |\phi_j\rangle\langle\phi_j|$ is an eigenvalue decomposition:

$$U^t U = I, \text{ i.e. } U \text{ is an isometry}$$

(ii) general case: $U = V \cdot W^t$, $V^t V = W^t W = I$, i.e.,

U is a partial isometry

(i.e. $U^t U$, $U U^t$ are projections)

Proof: We will first prove case (i).

" \Leftarrow ": Let $\sqrt{p_i} |\psi_i\rangle = \sum_j u_{ij} \sqrt{q_j} |\phi_j\rangle$.

$$\text{Then } \sum_i p_i |\psi_i\rangle\langle\psi_i| = \sum_i \left(\sum_j u_{ij} \sqrt{q_j} |\phi_j\rangle \right) \left(\sum_{j'} \overline{u_{ij'}} \sqrt{q_{j'}} \langle\phi_{j'}| \right)$$

$$= \sum_{j,j'} \sqrt{q_j} |\phi_j\rangle\langle\phi_{j'}| \sqrt{q_{j'}} \underbrace{\left(\sum_i \overline{u_{ij'}} u_{ij} \right)}$$

$$= \sum_j q_j |\phi_j\rangle\langle\phi_j|. \quad (u^t u)_{jj} = \delta_{jj}$$

" \Rightarrow ": First, assume $|\phi_j\rangle$ is an eigenbasis of P ,
and that all $g_j \neq 0$.

$$\text{Define } u_{ij} = \langle \phi_j | \psi_i \rangle \frac{\sqrt{p_i}}{\sqrt{g_j}}.$$

$$\begin{aligned} \text{Then, } \sum_j u_{ij} \sqrt{g_j} |\phi_j\rangle &= \sum_j \sqrt{g_j} |\phi_j\rangle \langle \phi_j | \psi_i \rangle \frac{\sqrt{p_i}}{\sqrt{g_j}} \\ &= \sqrt{p_i} |\psi_i\rangle, \end{aligned}$$

$$\begin{aligned} \text{and } \sum_i u_{ij} \overline{u_{ij'}} &= \sum_i \langle \phi_j | \psi_i \rangle \langle \psi_i | \phi_{j'} \rangle \frac{p_i}{\sqrt{g_j g_{j'}}} \\ &= \underbrace{\langle \phi_j | \underbrace{\sum_i p_i |\psi_i\rangle \langle \psi_i|}_{P} | \phi_{j'} \rangle}_{= g_j \delta_{jj'}} \frac{1}{\sqrt{g_j g_{j'}}} = \delta_{jj'} \end{aligned}$$

$\Rightarrow (u_{ij})$ is an isometry. □

General case: First, restrict to $\text{supp}(P)$, since all $|\phi_j\rangle, |\psi_i\rangle \in \text{supp}(P)$; then, all $g_i, p_j \neq 0$. Then, relate

$$\sum p_i |\psi_i\rangle \langle \psi_i| \xleftrightarrow{v_{ik}} \underbrace{\sum r_k |e_k\rangle \langle e_k|}_{\text{eigenbasis}} \xleftrightarrow{w_{jk}} \sum g_j |\phi_j\rangle \langle \phi_j|$$

& combine the isometries v_{ik} & w_{jk}

→ Homework. TMD

e) Unitary evolution & projective measurement for mixed states

How does a mixed state evolve under a unitary U ?

- Can be assessed in diff. ways, e.g. through purifications (here), or ensemble interpretation, or "Heisenberg picture" (= evolving meas. operators).

Consider state ρ & unitary U .

Let $|\psi\rangle = |\psi\rangle_{AB}$ be a purification of ρ ,

$$\text{tr}_B | \psi \rangle \langle \psi | = \rho_A.$$

$$\text{Then, } | \psi \rangle \longmapsto (U_A \otimes I_B) | \psi \rangle$$

$$\Rightarrow \rho_A = \text{tr}_B | \psi \rangle \langle \psi |$$

$$\longmapsto \text{tr}_B \left[(U_A \otimes I_B) | \psi \rangle \langle \psi | (U_A^\dagger \otimes I_B) \right]$$

$$= U_A \text{tr}_B \left[(I_A \otimes I_B) | \psi \rangle \langle \psi | (I_A \otimes I_B) \right] U_A^\dagger$$

$$= U_A \rho_A U_A^\dagger.$$

How does proj. measurement $\{E_u\}$ act on ρ_A ?

By construction of ρ_A , $p_u = \text{tr} [E_u \rho_A]$.

Post-meas. state:

$$\rho_{A,u} = \frac{1}{p_u} \text{tr}_B \left[(E_u \otimes I) | \psi \rangle \langle \psi | (E_u^\dagger \otimes I) \right]$$

$$= \frac{1}{p_u} E_u \rho_A E_u^\dagger.$$

(Note: Both derivations indep. of chosen purification
 \rightarrow well-defined.)